

## i-gel Psychomotor Competency

Candidate: \_\_\_\_\_

Agency: \_\_\_\_\_

Procedure	Successful	Unsuccessful
Takes appropriate PPE precautions		
Preoxygenates the patient appropriately		
Opens the i-gel package and removes the i-gel from the cradle		
Applies lubricant to the cradle		
Applies lubricant to the sides, back, and tip of the i-gel cuff		
Ensures that there is no lubricant in the bowl		
Replaces the i-gel in the cradle until ready to insert		
Removes the i-gel from the cradle and grasps it by the integral bite block		
Inserts the i-gel into the oropharynx with gentle constant pressure		
Stops insertion when definitive stop is felt		
Notes the placement of the i-gel at the teeth		
Applies waveform capnography and notes appropriate reading		
Listens to sounds over the epigastrium		
Listens to sounds in bilateral apexes		
Listens to sounds in bilateral bases		
Listens to sounds in bilateral midaxillaries		
Secures the device		
Ensures proper ventilatory rate and depth		
Recognizes need to remove the device		
Verbalizes continued care and reassessment		
Verbalizes need to document procedure in PCR		

### Critical Criteria

- \_\_\_\_\_ Failure to take or verbalize appropriate PPE precautions
- \_\_\_\_\_ Undue amount of force used to insert the i-gel
- \_\_\_\_\_ Candidate hyperventilated the patient pre/post i-gel insertion
- \_\_\_\_\_ Failure to check lung / epigastric sounds or to attach waveform capnography
- \_\_\_\_\_ Failure to secure the device properly

Evaluator Name \_\_\_\_\_ Credentials \_\_\_\_\_

Evaluator Signature \_\_\_\_\_ Date \_\_\_\_\_

Candidate Signature \_\_\_\_\_